TGU-29
UWThPh-2002-6
ZU-TH 7/02
hep-ph/0204071

arXiv:hep-ph/0204071v1 5 Apr 2002

Impact of CP phases on the search for sleptons $\tilde{\tau}$ and $\tilde{\nu}_\tau$

A. Bartl[a], K. Hidaka[b], T. Kernreiter[a] and W. Porod[c]

[a] *Institut für Theoretische Physik, Universität Wien, A-1090 Vienna, Austria*
[b] *Department of Physics, Tokyo Gakugei University, Koganei, Tokyo 184–8501, Japan*
[c] *Institut für Theoretische Physik, Universität Zürich, CH-8057 Zürich, Switzerland*

Abstract

We study the decays of the τ-sleptons ($\tilde{\tau}_{1,2}$) and τ-sneutrino ($\tilde{\nu}_\tau$) in the Minimal Supersymmetric Standard Model (MSSM) with complex parameters A_τ, μ and M_1 (U(1) gaugino mass). We show that the effect of the CP phases of these parameters on the branching ratios of $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ decays can be quite strong in a large region of the MSSM parameter space. This could have an important impact on the search for $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ and the determination of the MSSM parameters at future colliders.

So far most phenomenological studies on supersymmetric (SUSY) particle searches have been performed in the Minimal Supersymmetric Standard Model (MSSM) [1] with real SUSY parameters. Studies of the 3rd generation sfermions are particularly interesting because of the effects of the large Yukawa couplings. The lighter sfermion mass eigenstates may be relatively light and they could be thoroughly studied at an e^-e^+ linear collider [2]. They could also be copiously produced in the decays of heavier SUSY particles. An analysis of the decays of the 3rd generation sleptons $\tilde{\tau}_2$ and $\tilde{\nu}_\tau$ in the MSSM with real parameters was performed in Ref.[3], and phenomenological studies of production and decays of the 3rd generation sfermions at future e^-e^+ colliders in Ref.[4]. The assumption that all SUSY parameters are real, however, may be too restrictive. The higgsino mass parameter μ and the trilinear scalar coupling parameters A_f of the sfermions $\tilde{f}$ may be complex. In minimal Supergravity-type models the phase of μ (φ_μ) turns out to be restricted by the experimental data on electron and neutron electric dipole moments (EDMs) to a range $|\varphi_\mu| \lesssim 0.1 - 0.2$ for a universal scalar mass parameter $M_0 \lesssim 400$GeV, while the phase of the universal trilinear scalar coupling parameter A_0 is correlated with φ_μ, but otherwise unrestricted [5]. In more general models the phases of the parameters A_f of the 3rd generation sfermions are not restricted at one-loop level by the EDM data. However, there may be restrictions at two-loop level [6]. In a complete phenomenological analysis of production and decays of the SUSY particles one has to take into account that the μ and A_f may be complex. Furthermore, explicit CP violation in the Higgs sector can be induced by loop effects involving CP-violating interactions of Higgs bosons to top and bottom squark ($\tilde{t}$ and $\tilde{b}$) sector with complex parameters [7, 8]. It is found [9] that such effects of the complex phases on the phenomenology of the Higgs boson search could be quite significant. In principle, the imaginary parts of the possible complex SUSY parameters involved could most directly and unambiguously be determined by measuring relevant CP-violating observables; e.g. such analyses in τ-slepton ($\tilde{\tau}$) pair production in e^+e^- and $\mu^+\mu^-$ colliders were performed in Ref.[10].
On the other hand, the CP-conserving observables also can depend on the phases of the complex parameters because in general the mass-eigenvalues and the couplings of the SUSY particles (sparticles) involved are functions of the underlying complex parameters. For example, the branching ratios of the Higgs boson decays depend strongly on the complex phases of the $\tilde{t}$ and $\tilde{b}$ sectors [11].

In this article we study the effects of the complex phases of the stau and gaugino-higgsino sectors on the decay branching ratios of the staus $\tilde{\tau}_{1,2}$ and τ-sneutrino $\tilde{\nu}_\tau$ with $\tilde{\tau}_1(\tilde{\tau}_2)$ being the lighter (heavier) stau. We point out that these effects can be quite strong in a large region of the MSSM parameter space. This could have an important impact on the search for $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ and the determination of the MSSM parameters at future colliders.

First we summarize the MSSM parameters in our analysis. In the MSSM the stau sector is specified by the mass matrix in the basis $(\tilde{\tau}_L, \tilde{\tau}_R)$ [12]

$$\mathcal{M}^2_{\tilde{\tau}} = \begin{pmatrix} m^2_{\tilde{\tau}_L} & a^*_\tau m_\tau \\ a_\tau m_\tau & m^2_{\tilde{\tau}_R} \end{pmatrix} \tag{1}$$

with

$$\begin{aligned} m^2_{\tilde{\tau}_L} &= M^2_{\tilde{L}} + m^2_Z \cos 2\beta \, (\sin^2\theta_W - \frac{1}{2}) + m^2_\tau, & (2) \\ m^2_{\tilde{\tau}_R} &= M^2_{\tilde{E}} - m^2_Z \cos 2\beta \, \sin^2\theta_W + m^2_\tau, & (3) \\ a_\tau m_\tau &= (A_\tau - \mu^* \tan\beta)\, m_\tau \;=\; |a_\tau m_\tau|\, e^{i\varphi_{\tilde{\tau}}} \;(-\pi < \varphi_{\tilde{\tau}} \le \pi). & (4) \end{aligned}$$

$M_{\tilde{L},\tilde{E}}$ and A_τ are soft SUSY–breaking parameters, μ is the higgsino mass parameter, and $\tan\beta = v_2/v_1$ with v_1 (v_2) being the vacuum expectation value of the Higgs field H^0_1 (H^0_2). As the relative phase ξ between v_1 and v_2 is irrelevant in our analysis, we adopt the $\xi = 0$ scheme [8]. We take A_τ and μ as complex parameters: $A_\tau = |A_\tau|\, e^{i\varphi_{A_\tau}}$ and $\mu = |\mu|\, e^{i\varphi_\mu}$ with $-\pi < \varphi_{A_\tau,\mu} \le \pi$. Diagonalizing the matrix (1) one gets the mass eigenstates $\tilde{\tau}_1$ and $\tilde{\tau}_2$

$$\begin{pmatrix} \tilde{\tau}_1 \\ \tilde{\tau}_2 \end{pmatrix} = R^{\tilde{\tau}} \begin{pmatrix} \tilde{\tau}_L \\ \tilde{\tau}_R \end{pmatrix} = \begin{pmatrix} e^{i\varphi_{\tilde{\tau}}} \cos\theta_{\tilde{\tau}} & \sin\theta_{\tilde{\tau}} \\ -\sin\theta_{\tilde{\tau}} & e^{-i\varphi_{\tilde{\tau}}} \cos\theta_{\tilde{\tau}} \end{pmatrix} \begin{pmatrix} \tilde{\tau}_L \\ \tilde{\tau}_R \end{pmatrix} \tag{5}$$

with the masses $m_{\tilde{\tau}_1}$ and $m_{\tilde{\tau}_2}$ ($m_{\tilde{\tau}_1} < m_{\tilde{\tau}_2}$), and the mixing angle $\theta_{\tilde{\tau}}$

$$\begin{aligned} m^2_{\tilde{\tau}_{1,2}} &= \frac{1}{2}(m^2_{\tilde{\tau}_L} + m^2_{\tilde{\tau}_R} \mp \sqrt{(m^2_{\tilde{\tau}_L} - m^2_{\tilde{\tau}_R})^2 + 4|a_\tau m_\tau|^2}), & (6) \\ \theta_{\tilde{\tau}} &= \tan^{-1}(|a_\tau m_\tau|/(m^2_{\tilde{\tau}_1} - m^2_{\tilde{\tau}_R})) \quad (-\pi/2 \le \theta_{\tilde{\tau}} \le 0). & (7) \end{aligned}$$

The $\tilde{\tau}_L - \tilde{\tau}_R$ mixing is large if $|m^2_{\tilde{\tau}_L} - m^2_{\tilde{\tau}_R}| \lesssim |a_\tau m_\tau|$, which may be the case for large $\tan\beta$ and $|\mu|$. From Eqs.(6) and (7) we see that $m^2_{\tilde{\tau}_{1,2}}$ and $\theta_{\tilde{\tau}}$ depend on the phases only through a term $\cos(\varphi_{A_\tau} + \varphi_\mu)$. This phase dependence is strongest if $|A_\tau| \simeq |\mu| \tan\beta$. The mass of $\tilde{\nu}_\tau$ is given by

$$m^2_{\tilde{\nu}_\tau} = M^2_{\tilde{L}} + \frac{1}{2} m^2_Z \cos 2\beta. \tag{8}$$

The properties of charginos $\tilde{\chi}^\pm_i$ ($i = 1, 2$; $m_{\tilde{\chi}^\pm_1} < m_{\tilde{\chi}^\pm_2}$) and neutralinos $\tilde{\chi}^0_j$ ($j = 1, ..., 4$; $m_{\tilde{\chi}^0_1} < ... < m_{\tilde{\chi}^0_4}$) are determined by the parameters M_2, M_1, μ and $\tan\beta$, where M_2 and M_1 are the SU(2) and U(1) gaugino masses, respectively. We assume that M_2 and the gluino mass $m_{\tilde{g}}$ are real and that M_1 is complex: $M_1 = |M_1| e^{i\varphi_1}\,(-\pi < \varphi_1 \le \pi)$. Inspired by the gaugino mass unification we take $|M_1| = (5/3)\tan^2\theta_W M_2$ and $m_{\tilde{g}} = (\alpha_s(m_{\tilde{g}})/\alpha_2) M_2$. In the MSSM Higgs sector with explicit CP violation the mass-eigenvalues and couplings of the neutral and charged Higgs bosons H^0_1, H^0_2, H^0_3 ($m_{H^0_1} < m_{H^0_2} < m_{H^0_3}$) and $H^\pm$, including Yukawa and QCD radiative corrections, are fixed by $m_{H^+}, \tan\beta, \mu, m_t, m_b, M_{\tilde{Q}}, M_{\tilde{U}}, M_{\tilde{D}}, A_t, A_b, |M_1|, M_2$, and $m_{\tilde{g}}$ [8]. Here $M_{\tilde{Q},\tilde{U},\tilde{D}}$ and $A_{t,b}$ are the soft SUSY-breaking parameters in the $\tilde{t}$ and $\tilde{b}$ sectors, and $A_{t,b}$ are in general also complex: $A_{t,b} = |A_{t,b}| e^{i\varphi_{A_{t,b}}}$ $(-\pi < \varphi_{A_{t,b}} \le \pi)$. The neutral Higgs mass eigenstates H^0_1, H^0_2

and H_3^0 are mixtures of CP-even and CP-odd states ($\phi_{1,2}$ and a) due to the explicit CP violation in the Higgs sector. For the radiatively corrected masses and mixings of the Higgs bosons we use the formulae of Ref.[8]. We treat $M_{\{\tilde{L},\tilde{E},\tilde{Q},\tilde{U},\tilde{D}\}}$ and $A_{\{\tau,t,b\}}$ as free parameters since the ratios among them are highly model-dependent.

Here we list possible important decay modes of $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$:

$$\tilde{\tau}_1 \;\rightarrow\; \tau\,\tilde{\chi}_i^0\,,\;\nu_\tau\,\tilde{\chi}_j^- \tag{9}$$

$$\tilde{\tau}_2 \;\rightarrow\; \tau\,\tilde{\chi}_i^0\,,\;\nu_\tau\,\tilde{\chi}_j^-\,,\;\tilde{\tau}_1 Z^0\,,\;\tilde{\nu}_\tau W^-\,,\;\tilde{\tau}_1 H_k^0\,,\;\tilde{\nu}_\tau H^- \tag{10}$$

$$\tilde{\nu}_\tau \;\rightarrow\; \nu_\tau\,\tilde{\chi}_i^0\,,\;\tau\,\tilde{\chi}_j^+\,,\;\tilde{\tau}_1 W^+\,,\;\tilde{\tau}_1 H^+. \tag{11}$$

The decays into a gauge or Higgs boson in (10) and (11) are possible in case the mass splitting between the sleptons is sufficiently large [3]. The explicit expressions of the widths of the decays (9)-(11) in case of real SUSY parameters are given in [13]. Those for complex parameters can be obtained by using the corresponding masses and couplings (mixings) from Refs.[8, 12] and will be presented elsewhere [14].

The phase dependence of the widths stems from that of the involved mass-eigenvalues, mixings and couplings among the interaction-eigenfields. Here we summarize the latter phase dependence:

(I) $\tilde{\tau}_i$ sector:

- (a) $m_{\tilde{\tau}_{1,2}}$ are insensitive to the phases $(\varphi_{A_\tau}, \varphi_\mu)$ if the $\tilde{\tau}$-mixing term $|a_\tau m_\tau| \ll m_{\tilde{\tau}_L}^2 + m_{\tilde{\tau}_R}^2$. In most cases this is naturally fulfilled, because m_τ is small.
- (b) The $\tilde{\tau}$-mixing angle $\theta_{\tilde{\tau}}$ (given by $\tan 2\theta_{\tilde{\tau}} = 2|a_\tau m_\tau|/(m_{\tilde{\tau}_L}^2 - m_{\tilde{\tau}_R}^2)$) is sensitive to $(\varphi_{A_\tau}, \varphi_\mu)$ (via $\cos(\varphi_{A_\tau} + \varphi_\mu)$) if and only if $|m_{\tilde{\tau}_L}^2 - m_{\tilde{\tau}_R}^2|$ is small *and* $|A_\tau| \sim |\mu| \tan\beta$. Here note that $|m_{\tilde{\tau}_L}^2 - m_{\tilde{\tau}_R}^2| \simeq |M_{\tilde{L}}^2 - M_{\tilde{E}}^2| \simeq |m_{\tilde{\tau}_1}^2 - m_{\tilde{\tau}_2}^2|$ due to the smallness of $|a_\tau m_\tau|$.

(II) $\tilde{\chi}_i^0$ and $\tilde{\chi}_j^\pm$ sectors:

- (a) $m_{\tilde{\chi}_i^0}$ (i=1,...,4) and the $\tilde{\chi}^0$-mixing matrix are sensitive [insensitive] to the phases (φ_1, φ_μ) for small [large] $\tan\beta$.
- (b) $m_{\tilde{\chi}_{1,2}^\pm}$ and the $\tilde{\chi}^\pm$-mixing matrices are sensitive [insensitive] to φ_μ for small [large] $\tan\beta$.

(III) $\tilde{\nu}_\tau$ and $H^\pm$ sectors:
These sectors are independent of the phases.

(IV) H_i^0 sector:

- (a) $m_{H_i^0}$ (i=1,2,3) are sensitive [insensitive] to the phase sums $\varphi_{A_{t,b}} + \varphi_\mu$ for small [large] $\tan\beta$ [8].

(b) In general the H_i^0-mixing matrix (a real orthogonal 3×3 matrix O_{ij}) is sensitive to $\varphi_{A_{t,b}} + \varphi_\mu$ for any $\tan\beta$ [8].

(V) The couplings among the interaction-eigenfields:

(a) For the decays into fermions and gauge bosons in Eqs.(9)-(11), they are gauge couplings and/or tau Yukawa coupling (h_τ), and are independent of the phases.

(b) For the decays into Higgs bosons in Eqs.(10) and (11), the slepton-"chirality" flip (nonflip) couplings are dependent on (independent of) the phases φ_{A_τ} and φ_μ;

$$\begin{aligned} C(\tilde\nu_\tau^\dagger\tilde\tau_R H^+) &\sim \cos\beta\, h_\tau(A_\tau^*\tan\beta+\mu) && (12)\\ C(\tilde\tau_L^\dagger\tilde\tau_R\phi_1) &\sim h_\tau A_\tau^* && (13)\\ C(\tilde\tau_L^\dagger\tilde\tau_R\phi_2) &\sim h_\tau\mu && (14)\\ C(\tilde\tau_L^\dagger\tilde\tau_R a) &\sim \cos\beta\, h_\tau(A_\tau^*\tan\beta+\mu) && (15)\\ C(\tilde\nu_\tau^\dagger\tilde\tau_L H^+) &\sim C(\tilde\tau_L^\dagger\tilde\tau_L\phi_{1,2}) \sim C(\tilde\tau_R^\dagger\tilde\tau_R\phi_{1,2}) \sim g m_W && (16) \end{aligned}$$

with

$$h_\tau = g m_\tau/(\sqrt{2} m_W\cos\beta). \qquad (17)$$

Here $\phi_i = O_{ij}H_j^0$ (i=1,2) and $a = O_{3j}H_j^0$ are the CP-even and CP-odd neutral Higgs bosons, respectively [8].

From the facts (I)-(V) the widths (and hence the branching ratios) of the decays (9)-(11) are expected to be sensitive to the phases (φ_{A_τ}, φ_μ, φ_1, $\varphi_{A_{t,b}}+\varphi_\mu$) in a large region of the MSSM parameter space.

Now we turn to the numerical analysis of the $\tilde\tau_{1,2}$ and $\tilde\nu_\tau$ decay branching ratios. We calculate the widths of all possibly important two-body decay modes of Eqs.(9)-(11). Three-body decays are negligible in this study. We take $m_\tau = 1.78$ GeV, $m_t = 175$ GeV, $m_b = 5$ GeV, $m_Z = 91.2$ GeV, $\sin^2\theta_W = 0.23$, $m_W = m_Z\cos\theta_W$, $\alpha(m_Z) = 1/129$, and $\alpha_s(m_Z) = 0.12$ (with $\alpha_s(Q) = 12\pi/((33-2n_f)\ln(Q^2/\Lambda_{n_f}^2))$ for the determination of $m_{\tilde g}(=(\alpha_s(m_{\tilde g})/\alpha_2)M_2)$, n_f being the number of quark flavors), where $m_{t,b}$ are pole masses of t and b quarks. In order not to vary too many parameters we fix $m_{\tilde\tau_1} = 240$GeV, $m_{H^+} = 180$GeV, $M_{\tilde Q} = M_{\tilde U} = M_{\tilde D} = |A_t| = |A_b| = 1000$GeV, and $\varphi_{A_t} = \varphi_{A_b} = 0$ for simplicity. In our numerical study we take $\tan\beta$, M_2, $m_{\tilde\tau_2}$, $|A_\tau|$, $|\mu|$, φ_{A_τ}, φ_μ and φ_1 as input parameters. Note that for a given set of the input parameters we have two solutions for ($M_{\tilde L}$, $M_{\tilde E}$) corresponding to the two cases $m_{\tilde\tau_L} \geq m_{\tilde\tau_R}$ and $m_{\tilde\tau_L} < m_{\tilde\tau_R}$. In the plots we impose the following conditions in order to respect experimental and theoretical constraints:

(i) $m_{\tilde\chi_1^\pm} > 103$ GeV, $m_{\tilde\chi_1^0} > 50$ GeV, $m_{\tilde\tau_1,\tilde t_1,\tilde b_1} > 100$ GeV, $m_{\tilde\tau_1,\tilde t_1,\tilde b_1} > m_{\tilde\chi_1^0}$, $m_{H_1^0} > 110$ GeV,

(ii) $|A_\tau|^2 < 3\,(M_{\tilde{L}}^2 + M_{\tilde{E}}^2 + m_1^2)$, $|A_t|^2 < 3\,(M_{\tilde{Q}}^2 + M_{\tilde{U}}^2 + m_2^2)$, and $|A_b|^2 < 3\,(M_{\tilde{Q}}^2 + M_{\tilde{D}}^2 + m_1^2)$, where $m_1^2 = (m_{H^+}^2 + m_Z^2 \sin^2\theta_W)\sin^2\beta - \frac{1}{2}\,m_Z^2$ and $m_2^2 = (m_{H^+}^2 + m_Z^2 \sin^2\theta_W)\cos^2\beta - \frac{1}{2}\,m_Z^2$,

(iii) $\Delta\rho\,(\tilde{t}-\tilde{b}) < 0.0012$ [15] using the formula of [16].

Condition (i) is imposed to satisfy the experimental mass bounds from LEP [17]. (ii) is the approximate necessary condition for the tree-level vacuum stability [18]. (iii) constrains μ and $\tan\beta$ (in the squark sector). We do not impose the $b \to s\gamma$ constraint [19] since it strongly depends on the details of the flavour structures of the squarks, including the generation-mixing.

In general the experimental upper limits on the electron and neutron electric dipole moments (EDMs) strongly constrain the SUSY CP phases [5]. One interesting possibility for evading these constraints is to invoke large masses (much above the TeV scale) for the first two generations of the sfermions [20], keeping the third generation sfermions relatively light ($\lesssim$ 1 TeV). In such a scenario (φ_1, φ_μ) and the CP phases in the third generation (φ_{A_τ}, φ_{A_t}, φ_{A_b}) are practically unconstrained [20]. We take this scenario. The deviation of the recent data on the muon g-2 from the Standard Model prediction is no longer significant [21], which allows of our scenario. We have checked that the electron and neutron EDM constraints at two-loop level [6] are fulfilled in the numerical examples studied in this article.

In Fig.1 we plot the contours of the branching ratios of the $\tilde{\tau}_1$ decays $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_1^0)$ and $B(\tilde{\tau}_1 \to \nu_\tau\tilde{\chi}_1^-)$ in the $|A_\tau| - |\mu|$ plane for $\tan\beta = 3$, $M_2 = 200$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 250$GeV, $\varphi_1 = 0$, and (φ_{A_τ}, φ_μ) = (0,0), (π/2,0), (0,π/2) in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$; in the case $m_{\tilde{\tau}_L} \geq m_{\tilde{\tau}_R}$ we have obtained similar results. As expected, these branching ratios are very sensitive to the phases φ_{A_τ} and φ_μ in a sizable region of the $|A_\tau| - |\mu|$ plane. As can be seen from item (I), in this case the $\tilde{\tau}$-mixing angle $\theta_{\tilde{\tau}}$ is sensitive to $\cos(\varphi_{A_\tau} + \varphi_\mu)$ for $|A_\tau| \sim 3|\mu|$, which is the main cause for the difference between the case of (φ_{A_τ}, φ_μ) = (0,0) and those of (π/2,0) and (0,π/2). Furthermore from item (II) one sees that the masses and mixing-matrices of the $\tilde{\chi}_i^0$ and $\tilde{\chi}_j^\pm$ are sensitive to φ_μ, which is the main reason for the difference between the case of (φ_{A_τ}, φ_μ) = (0,π/2) and those of (0,0) and (π/2,0). Here note also item (V)-(a).

In Fig.2 we plot the contours of the $\tilde{\tau}_1$ decay branching ratios $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_1^0)$, $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_2^0)$, and $B(\tilde{\tau}_1 \to \nu_\tau\tilde{\chi}_1^-)$ in the $\varphi_{A_\tau} - \varphi_\mu$ plane for $\tan\beta = 3$, $M_2 = 200$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 255$GeV, $|A_\tau| = 600$GeV, $|\mu| = 350$GeV, and $\varphi_1 = 0$ in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$; in the case $m_{\tilde{\tau}_L} \geq m_{\tilde{\tau}_R}$ we have obtained similar results. One sees that these branching ratios depend on the CP phases φ_{A_τ} and φ_μ quite strongly, as expected from items (I) and (II).

In Fig.3 we show the φ_μ dependence of the $\tilde{\tau}_1$ decay branching ratios for $\tan\beta = 3, 30$ (with $\varphi_1 = 0$) and $\varphi_1 = 0, \pi/2$ (with $\tan\beta = 3$) with $M_2 = 300$GeV, $m_{\tilde{\tau}_2} = 500$GeV, $|A_\tau| = 600$GeV, $|\mu| = 200$GeV, and $\varphi_{A_\tau} = 0$ in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$; in the case $m_{\tilde{\tau}_L} \geq m_{\tilde{\tau}_R}$

we have obtained similar results. From Fig.3a we see that the $\tilde{\tau}_1$ decay branching ratios are sensitive (insensitive) to φ_μ for small (large) $\tan\beta$ as is expected from items (I), (II), and (V)-(a). In general the $\tilde{\tau}_1$ decay branching ratios are insensitive to all of the phases (φ_{A_τ}, φ_μ, φ_1) for large $\tan\beta$ according to items (I), (II) and (V)-(a). From Fig.3b we find that they are sensitive to φ_μ and φ_1 for small $\tan\beta(=3)$ where the masses and mixings of $\tilde{\chi}^0_i$ [$\tilde{\chi}^\pm_j$] are sensitive to (φ_μ, φ_1) [φ_μ] (though $\theta_{\tilde{\tau}}$ is insensitive to φ_μ in this case) as seen from items (I) and (II).

In Fig.4 we show the φ_{A_τ} dependence of the $\tilde{\tau}_2$ decay branching ratios for $\varphi_\mu = 0$ and $\pi/2$ in the two cases of small (a) and large (b,c) mass difference $|m_{\tilde{\tau}_1} - m_{\tilde{\tau}_2}|$. For the former case we take ($m_{\tilde{\tau}_1}$(GeV), $m_{\tilde{\tau}_2}$(GeV), $\tan\beta$, M_2(GeV), $|A_\tau|$(GeV), $|\mu|$(GeV), φ_1) = (240, 260, 6, 500, 600, 150, 0) with $m_{\tilde{\tau}_L} \geq m_{\tilde{\tau}_R}$, whereas for the latter case we take ($m_{\tilde{\tau}_1}$(GeV), $m_{\tilde{\tau}_2}$(GeV), $\tan\beta$, M_2(GeV), $|A_\tau|$(GeV), $|\mu|$(GeV), φ_1) = (240, 500, 30, 400, 900, 800, 0) with $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$. In Fig.4a (where $|m_{\tilde{\tau}_1} - m_{\tilde{\tau}_2}|$ and $|m_{\tilde{\nu}_\tau} - m_{\tilde{\tau}_2}|$ are so small that the bosonic decays in Eq.(10) are kinematically forbidden) we see that the $\tilde{\tau}_2$ decay branching ratios are very sensitive to φ_{A_τ} and φ_μ as expected from items (I) and (II). In Figs.4b and 4c (where $|m_{\tilde{\tau}_1} - m_{\tilde{\tau}_2}|(\sim |m_{\tilde{\nu}_\tau} - m_{\tilde{\tau}_2}|)$ is so large that the bosonic decays in Eq.(10) also are allowed) we find that the branching ratios of the Higgs boson modes are rather sensitive to φ_{A_τ} and φ_μ as expected from items (IV)-(b) and (V)-(b); here note that in this case $(\tilde{\tau}_1, \tilde{\tau}_2) \sim (\tilde{\tau}_L, \tilde{\tau}_R)$ due to the smallness of the $\tilde{\tau}_L$-$\tilde{\tau}_R$ mixing term and hence that the bosonic decays of $\tilde{\tau}_2$ are basically the decays of $\tilde{\tau}_R$ into $(\tilde{\tau}_L, \tilde{\nu}_\tau)$.

In Fig.5 we show the φ_μ dependence of the $\tilde{\nu}_\tau$ decay branching ratios for $\varphi_1 = 0$ and $\pi/2$ in the case of $\tan\beta = 3$, $M_2 = 500$GeV, $m_{\tilde{\tau}_2} = 500$GeV, $|A_\tau| = 600$GeV, $|\mu| = 150$GeV, $\varphi_{A_\tau} = 0$, and $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$ (for which $m_{\tilde{\nu}_\tau} \sim m_{\tilde{\tau}_L} \sim m_{\tilde{\tau}_1}$ and hence the bosonic decays in Eq.(11) are kinematically forbidden). We see that the $\tilde{\nu}_\tau$ decay branching ratios are quite sensitive to φ_μ and φ_1 as expected from item (II). In general the $\tilde{\nu}_\tau$ decay branching ratios are insensitive to all of the phases (φ_{A_τ}, φ_μ, φ_1) for large $\tan\beta$ as can be seen from items (I)∼(III) and (V). Furthermore, they tend to be insensitive to φ_{A_τ} for any $\tan\beta$ as can be seen from items (I)∼(III) and (V), except for some special cases such as the case with $m_{\tilde{\tau}_L} > m_{\tilde{\tau}_R}$, $|m_{\tilde{\tau}_1} - m_{\tilde{\tau}_2}| > m_{H^+}$, small $\tan\beta$, and $|\mu| \ll m_{\tilde{\nu}_\tau}(\sim m_{\tilde{\tau}_2}) < |M_{1,2}|$, where $m_{\tilde{\nu}_\tau} > m_{\tilde{\tau}_1} + m_{H^+}$, $m_{\{\tilde{\chi}^0_{1,2}, \tilde{\chi}^\pm_1\}} < m_{\tilde{\nu}_\tau} < m_{\{\tilde{\chi}^0_{3,4}, \tilde{\chi}^\pm_2\}}$, $\tilde{\tau}_1 \sim \tilde{\tau}_R$, and $(\tilde{\chi}^0_{1,2}, \tilde{\chi}^\pm_1)[(\tilde{\chi}^0_{3,4}, \tilde{\chi}^\pm_2)]$ are higgsino-like [gaugino-like]; in this case only the $\tau\tilde{\chi}^+_1$ and $\tilde{\tau}_1 H^+$ modes dominate the $\tilde{\nu}_\tau$ decay, and the coupling $C(\tilde{\nu}_\tau \tilde{\tau}^\dagger_1 H^-) \sim C(\tilde{\nu}_\tau \tilde{\tau}^\dagger_R H^-) \sim \cos\beta\, h_\tau (A_\tau \tan\beta + \mu^*)$ can be rather sensitive to φ_{A_τ} (though the $\theta_{\tilde{\tau}}$ is insensitive to φ_{A_τ}), which results in significant φ_{A_τ}-dependence of the $\tilde{\nu}_\tau$ decay branching ratios. We have checked that this can be easily realized indeed.

As for the τ-lepton EDM (d_τ), we have checked that in the MSSM parameter region considered here the predicted range of it is well below the current experimental limit ($|d_\tau| < 3.1 \times 10^{-16} e\, cm$) [22] and most likely also below the expected sensitivity of future experiments to measure this EDM: we find that in the parameter region considered here $|d_\tau| \lesssim 10^{-20} e\, cm$, which is obtained by using the corresponding formulas in Ref. [23] with m_e replaced by m_τ.

The CP phases can significantly affect not only CP-violating observables such as the lepton EDM but also CP-conserving quantities such as the branching ratios of the $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ decays. Hence the possible sizable phases could have important consequences for the determination of the fundamental MSSM parameters by measurements of CP-conserving observables from which they are extracted.

In conclusion, we have shown that the effect of the CP phases of the complex parameters A_τ, μ and M_1 on the branching ratios of the $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ decays can be quite strong in a large region of the MSSM parameter space. This could have an important impact on the search for $\tilde{\tau}_{1,2}$ and $\tilde{\nu}_\tau$ and the determination of the MSSM parameters at future colliders.

Acknowledgements

The authors thank A. Pilaftsis and C.E.M. Wagner for useful correspondence. This work was supported by the 'Fonds zur Förderung der wissenschaftlichen Forschung' of Austria, Project No. P13139-PHY, and by the European Community's Human Potential Programme under contract HPRN-CT-2000-00149. W.P. is supported by 'Fonds zur Förderung der wissenschaftlichen Forschung' of Austria, Erwin Schrödinger fellowship Nr. J2095, and partly by the 'Schweizer Nationalfonds'.

References

[1] For reviews, see:
H.P. Nilles, Phys. Rep. 110 (1984) 1; H.E. Haber and G.L. Kane, Phys. Rep. 117 (1985) 75; R. Barbieri, Riv. Nuov. Cim. 11 (1988) 1.

[2] E. Accomando et al., Phys. Rep. 229 (1998) 1.

[3] A. Bartl, H. Eberl, K. Hidaka, S. Kraml, T. Kon, W. Majerotto, W. Porod, and Y. Yamada, Phys. Lett. B 460 (1999) 157.

[4] M. M. Nojiri, Phys. Rev. D 51 (1995) 6281; M. M. Nojiri, K. Fujii and T. Tsukamoto, Phys. Rev. D 54 (1996) 6756; A. Bartl, H. Eberl, S. Kraml, W. Majerotto, W. Porod and A. Sopczak, Z. Phys. C 76 (1997) 549; A. Bartl, H. Eberl, S. Kraml, W. Majerotto and W. Porod, Eur. Phys. J. direct C 6 (2000) 1.

[5] P. Nath, Talk at The 9th International Conference on Supersymmetry and Unification of Fundamental Interactions, 11-17 June 2001, Dubna (hep-ph/0107325) and references therein.

[6] D. Chang, W. Keung and A. Pilaftsis, Phys. Rev. Lett. 82 (1999) 900.

[7] A. Pilaftsis, Phys. Lett. B 435 (1998) 88; A. Pilaftsis and C.E.M. Wagner, Nucl. Phys. B553 (1999) 3; S.Y. Choi, M. Drees and J.S. Lee, Phys. Lett. B 481 (2000) 57; G.L. Kane and L.T. Wang, Phys. Lett. B 488 (2000) 383.

[8] M. Carena, J. Ellis, A. Pilaftsis and C.E.M. Wagner, Nucl. Phys. B586(2000)92.

[9] M. Carena, J. Ellis, A. Pilaftsis and C.E.M. Wagner, Phys. Lett. B 495 (2000) 155.

[10] S.Y. Choi and M. Drees, Phys. Lett. B 435 (1998) 356; S.Y. Choi and M. Drees, Phys. Rev. Lett. 81 (1998) 5509; S.Y. Choi, M. Drees, B. Gaissmaier and J.S. Lee, Phys. Rev. D 64 (2001) 095009.

[11] S.Y. Choi and J.S. Lee, Phys. Rev. D 61 (1999) 015003; S.Y.Choi, K. Hagiwara and J.S. Lee, Phys. Rev. D 64 (2001) 032004.

[12] J. Ellis and S. Rudaz, Phys. Lett. B 128 (1983) 248; J. F. Gunion and H. E. Haber, Nucl. Phys. B 272 (1986) 1; B 402 (1993) 567 (E).

[13] W. Porod, PhD Thesis, Univ. Vienna, 1997 (hep-ph/9804208).

[14] A. Bartl, K. Hidaka, T. Kernreiter and W. Porod, in preparation.

[15] G. Altarelli, R. Barbieri, and F. Caravaglios, Int. J. Mod. Phys. A 13 (1998) 1031.

[16] M. Drees and K. Hagiwara, Phys. Rev. D 42 (1990) 1709; S.K. Kang and J. D. Kim, Phys. Rev. D 62 (2000) 071901.

[17] G.G. Hanson, Talk at The 20th International Symposium on Lepton and Photon Interactions at High Energies, 23 - 28 July 2001, Rome (hep-ex/0111058).

[18] J. P. Derendinger and C. A. Savoy, Nucl. Phys. B 237 (1984) 307; J.A. Casas and S. Dimopoulos, Phys. Lett. B 387 (1996) 107.

[19] H. Baer, M. Brhlik, D. Castaño, and X. Tata, Phys. Rev. D 58 (1998) 015007, and references therein.

[20] A.G. Cohen, D.B. Kaplan and A.E. Nelson, Phys. Lett. B 388 (1996) 588; A.G. Akeroyd, Y.-Y. Keum and S. Recksiegel, Phys. Lett. B 507 (2001) 252 and references therein; Ref.[8] and references therein.

[21] M. Knecht and A. Nyffeler, hep-ph/0111058; M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, hep-ph/0111059; M. Hayakawa and T. Kinoshita, hep-ph/0112102.

[22] Review of Particle Physics, Euro. Phys. J. C 3 (1998) 286.

[23] A. Bartl, T. Gajdosik, W. Porod, P. Stockinger and H. Stremnitzer, Phys. Rev. D 60 (1999) 073003.

Figure Captions

Figure 1: Contours of the $\tilde{\tau}_1$ decay branching ratios $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_1^0)$ (a,b,c) and $B(\tilde{\tau}_1 \to \nu_\tau\tilde{\chi}_1^-)$ (d,e,f) in the $|A_\tau| - |\mu|$ plane for $\tan\beta = 3$, $M_2 = 200$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 250$GeV, $\varphi_1 = 0$, and $(\varphi_{A_\tau}, \varphi_\mu) = (0{,}0)$ (a,d), $(\pi/2{,}0)$ (b,e), and $(0{,}\pi/2)$ (c,f) in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$. The blank areas are excluded by the conditions (i) to (iii) given in the text and the inequality $(m_{\tilde{\tau}_L}^2 - m_{\tilde{\tau}_R}^2)^2 = (m_{\tilde{\tau}_1}^2 - m_{\tilde{\tau}_2}^2)^2 - (2|a_\tau m_\tau|)^2 \geq 0$. The excluded region is different in each plot mainly due to the fact that the inequality depends on the phases $(\varphi_{A_\tau}, \varphi_\mu)$.

Figure 2: Contours of the $\tilde{\tau}_1$ decay branching ratios $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_1^0)$ (a), $B(\tilde{\tau}_1 \to \tau\tilde{\chi}_2^0)$ (b), and $B(\tilde{\tau}_1 \to \nu_\tau\tilde{\chi}_1^-)$ (c) in the $\varphi_{A_\tau} - \varphi_\mu$ plane for $\tan\beta = 3$, $M_2 = 200$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 255$GeV, $|A_\tau| = 600$GeV, $|\mu| = 350$GeV, and $\varphi_1 = 0$ in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$.

Figure 3: φ_μ dependence of the $\tilde{\tau}_1$ decay branching ratios for $\tan\beta = 3, 30$ (with $\varphi_1 = 0$) (a) and $\varphi_1 = 0, \pi/2$ (with $\tan\beta = 3$) (b) with $M_2 = 300$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 500$GeV, $|A_\tau| = 600$GeV, $|\mu| = 200$GeV, and $\varphi_{A_\tau} = 0$ in the case $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$. In Fig.a the solid and dashed lines are for $\tan\beta = 3$ and 30, respectively. In Fig.b the solid and dashed lines are for $\varphi_1 = 0$ and $\pi/2$, respectively.

Figure 4: φ_{A_τ} dependence of the $\tilde{\tau}_2$ decay branching ratios for $\varphi_\mu = 0$ (solid lines) and $\pi/2$ (dashed lines) in two cases: (a) small $\tilde{\tau}$-mass-splitting case with $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 260$GeV, $\tan\beta = 6$, $M_2 = 500$GeV, $|A_\tau| = 600$GeV, $|\mu| = 150$GeV, $\varphi_1 = 0$, and $m_{\tilde{\tau}_L} \geq m_{\tilde{\tau}_R}$ (for which $m_{\tilde{\nu}_\tau} \sim m_{\tilde{\tau}_2}$), and (b,c) large $\tilde{\tau}$-mass-splitting case with $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 500$GeV, $\tan\beta = 30$, $M_2 = 400$GeV, $|A_\tau| = 900$GeV, $|\mu| = 800$GeV, $\varphi_1 = 0$, and $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$ (for which $m_{\tilde{\nu}_\tau} \sim m_{\tilde{\tau}_1}$). In the latter case (Figs.b and c) we have $(m_{H_1^0}, m_{H_2^0}, m_{H_3^0}) = (117.4, 159.2, 159.5)$(GeV) and $(117.7, 150.6, 151.2)$(GeV) for $\varphi_\mu = 0$ and $\pi/2$, respectively. The branching ratios $B(\tilde{\tau}_2 \to \tilde{\tau}_1 Z^0)\,(\sim 10\%)$ and $B(\tilde{\tau}_2 \to \tilde{\nu}_\tau W^-)\,(\sim 20\%)$, which are rather insensitive to φ_{A_τ}, are not shown in Figs.b and c.

Figure 5: φ_μ dependence of the $\tilde{\nu}_\tau$ decay branching ratios for $\varphi_1 = 0$ (solid lines) and $\pi/2$ (dashed lines) with $\tan\beta = 3$, $M_2 = 500$GeV, $m_{\tilde{\tau}_1} = 240$GeV, $m_{\tilde{\tau}_2} = 500$GeV, $|A_\tau| = 600$GeV, $|\mu| = 150$GeV, $\varphi_{A_\tau} = 0$, and $m_{\tilde{\tau}_L} < m_{\tilde{\tau}_R}$ (for which $m_{\tilde{\nu}_\tau} \sim m_{\tilde{\tau}_1}$).


(a)
(b)
(c)
(d)
(e)
(f)
|μ| (GeV)
|A_τ| (GeV)
0
100
200
300
400
500
600
700
0.1
0.2
0.3
0.4
0.5
0.6
0.25
0.35
0.45

Fig.1


(a)
0.6
0.5
0.4
0.3
0.22
0.3
0.4
0.5
0.6
(b)
0.15
0.2
0.25
0.28
0.2
0.15
(c)
0.25
0.3
0.35
0.4
0.45
0.5
0.45
0.4
0.35
0.4
0.3
0.25
150
100
50
0
-50
-100
-150
-150
-100
-50
0
50
100
150
φ_{μ} (deg)
$\varphi_{A\tau}$ (deg)

Fig.2


(a)
0.8
0.6
0.4
0.2
0.0
B($\tilde{\tau}_1$→X)
$\tau\,\tilde{\chi}_1^0$
$\tau\,\tilde{\chi}_2^0$
$\nu_\tau\,\tilde{\chi}_1^-$
-150
-100
-50
0
50
100
150
φ_μ (deg)
(b)
0.8
0.6
0.4
0.2
0.0
B($\tilde{\tau}_1$→X)
$\tau\,\tilde{\chi}_1^0$
$\tau\,\tilde{\chi}_2^0$
$\tau\,\tilde{\chi}_2^0$
$\nu_\tau\,\tilde{\chi}_1^-$
$\nu_\tau\,\tilde{\chi}_1^-$
-150
-100
-50
0
50
100
150
φ_μ (deg)

Fig.3

(a)
$B(\tilde{\tau}_2 \to X)$
φ_{A_τ} (deg)
$\tau\,\tilde{\chi}^0_2$
$\nu_\tau\,\tilde{\chi}^-_1$
$\tau\,\tilde{\chi}^0_1$
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
-150
-100
-50
0
50
100
150
(b)
$\varphi_\mu = 0$
$\tau\,\tilde{\chi}^0_1$
$\tilde{\nu}_\tau\,H^-$
$\tilde{\tau}_1\,H^0_2$
$\tilde{\tau}_1\,H^0_3$
$\tilde{\tau}_1\,H^0_1$
$\tilde{\tau}_1\,H^0_3$
$B(\tilde{\tau}_2 \to X)$
φ_{A_τ} (deg)
0.00
0.05
0.10
0.15
0.20
0.25
-150
-100
-50
0
50
100
150
(c)
$\varphi_\mu = \pi / 2$
$\tau\,\tilde{\chi}^0_1$
$\tilde{\nu}_\tau\,H^-$
$\tilde{\tau}_1\,H^0_3$
$\tilde{\tau}_1\,H^0_2$
$\tilde{\tau}_1\,H^0_1$
$B(\tilde{\tau}_2 \to X)$
φ_{A_τ} (deg)
0.00
0.05
0.10
0.15
0.20
0.25
-150
-100
-50
0
50
100
150

Fig.4


0.8
0.6
0.4
0.2
0.0
B($\tilde{\nu}_\tau$ → X)
-150
-100
-50
0
50
100
150
φ_μ (deg)
$\tau\ \tilde{\chi}_1^+$
$\nu_\tau\ \tilde{\chi}_1^0$
$\nu_\tau\ \tilde{\chi}_2^0$

Fig.5